

Oregon Grass Seed Crop Estimates¹

Estimates for 2016r, 2017r and 2018p

	Harvested Acres	Yield (lb/a)	Production (000 lbs.)	Price per cwt	Sales (000 \$)
2016 Annual ryegrass	115,490	1,881	217,237	33.00	71,688
2017 Annual ryegrass	120,250	1,721	206,909	42.00	86,902
2018 Annual ryegrass	123,780	1,791	221,712	49.00	108,639
2016 Perennial ryegrass	95,130	1,537	146,215	82.00	119,816
2017 Perennial ryegrass	83,450	1,476	123,208	79.00	97,334
2018 Perennial ryegrass	74,030	1,498	110,916	77.00	85,405
2016 Tall fescue	135,020	1,534	207,121	74.00	153,269
2017 Tall fescue	134,370	1,542	207,148	82.00	169,861
2018 Tall fescue	142,570	1,683	239,902	87.00	208,715
2016 Kentucky bluegrass	18,280	1,184	21,637	137.00	29,643
2017 Kentucky bluegrass	20,650	915	18,887	140.00	26,442
2018 Kentucky bluegrass	22,200	1,141	25,336	143.00	36,230
2016 Rough bluegrass	1,030	1,039	1,071	135.00	1,446
2017 Rough bluegrass	1,080	1,000	1,080	135.00	1,458
2018 Rough bluegrass	1,030	1,250	1,288	165.00	2,125
2016 Orchardgrass	13,330	923	12,300	212.00	26,076
2017 Orchardgrass	15,190	1,046	15,889	225.00	35,750
2018 Orchardgrass	16,170	1,005	16,250	203.00	32,988
2016 Chewings fescue	8,450	1,148	9,737	101.00	9,834
2017 Chewings fescue	8,790	1,141	10,029	105.00	10,531
2018 Chewings fescue	9,080	1,315	11,936	108.00	12,891
2016 Red fescue	9,160	1,047	9,588	101.00	9,684
2017 Red fescue	11,370	1,009	11,469	103.00	11,813
2018 Red fescue	10,810	1,115	12,055	107.00	12,899
2016 Hard fescue	2,470	1,093	2,699	140.00	3,779
2017 Hard fescue	2,280	1,009	2,300	140.00	3,220
2018 Hard fescue	2,710	1,414	3,831	130.00	4,980
2016 Colonial bentgrass	3,080	397	1,222	250.00	3,055
2017 Colonial bentgrass	3,030	379	1,147	250.00	2,868
2018 Colonial bentgrass	3,490	361	1,261	213.00	2,686
2016 Creeping bentgrass	3,860	589	2,274	340.00	7,732
2017 Creeping bentgrass	3,590	719	2,581	350.00	9,034
2018 Creeping bentgrass	3,540	762	2,698	365.00	9,848
2016 Total Grass	405,300		631,101		436,022
2017 Total Grass	404,050		600,647		455,213
2018 Total Grass	409,410		647,185		517,406

¹Data collected by the Nicole Anderson, Will Jessie, Ruijun Qin, Christy Tanner, Carol Tollefson, Bestsy Verhoeven and Darrin Walenta, Oregon State University, and compiled by Nicole Anderson, Extension Field Crops Agronomist, OSU.

r = revised, p = preliminary.