

# NATIVE WILDFLOWER SEED YIELD IN RESPONSE TO MODEST IRRIGATION

Clinton C. Shock, Erik B. G. Feibert, Alicia Rivera, and Kyle D. Wieland, Malheur Experiment Station, Oregon State University, Ontario, OR

Nancy Shaw and Francis Kilkenny, U.S. Forest Service, Rocky Mountain Research Station, Boise, ID

## Introduction

Commercial seed production of native wildflowers is necessary to provide the quantity of seed needed for restoration of Intermountain West rangelands. Native wildflower plants may not be well adapted to croplands. Native plants are often not competitive with crop weeds in cultivated fields, and this poor competitiveness with weeds could limit wildflower seed production. Both sprinkler and furrow irrigation could provide supplemental water for seed production, but these irrigation systems risk further encouraging weeds. Also, sprinkler and furrow irrigation can lead to the loss of plant stand and seed production due to fungal pathogens. By burying drip tape at a 12-inch depth and avoiding wetting the soil surface, we designed experiments to assure flowering and seed set without undue encouragement of weeds or opportunistic diseases. The trials reported here tested effects of three low rates of irrigation on seed yield of 14 native wildflower species (Table 1).

Table 1. Wildflower species planted at the Malheur Experiment Station, Oregon State University, Ontario, OR.

Species	Common name	Longevity	Row spacing (inches)
<i>Chaenactis douglasii</i>	Douglas' dustymaiden	perennial	30
<i>Crepis intermedia</i> <sup>a</sup>	limestone hawksbeard	perennial	30
<i>Cymopterus bipinnatus</i> <sup>b</sup>	Hayden's cymopterus	perennial	30
<i>Enceliopsis nudicaulis</i>	nakedstem sunray	perennial	30
<i>Heliomeris multiflora</i>	showy goldeneye	perennial	30
<i>Ipomopsis aggregata</i>	scarlet gilia	biennial	15
<i>Ligusticum canbyi</i>	Canby's licorice-root	perennial	30
<i>Ligusticum porteri</i>	Porter's licorice-root	perennial	30
<i>Machaeranthera canescens</i>	hoary tansyaster	perennial	30
<i>Nicotiana attenuata</i>	coyote tobacco	perennial	30
<i>Phacelia linearis</i>	threadleaf phacelia	annual	15
<i>Phacelia hastata</i>	silverleaf phacelia	perennial	15
<i>Thelypodium milleflorum</i>	manyflower thelypody	biennial	30
<i>Achillea millefolium</i>	common yarrow	perennial	30

<sup>a</sup>Planted in the fall of 2011.

<sup>b</sup>Recently classified as *Cymopterus nivalis* S. Watson "snowline springparsley." Planted in the fall of 2009.

## Materials and Methods

### Plant establishment

Each wildflower species was planted on 60-inch beds in rows 450 ft long on Nyssa silt loam at the Malheur Experiment Station, Ontario, Oregon. The soil had a pH of 8.3 and 1.1% organic matter. In October 2012, drip tape (T-Tape TSX 515-16-340) was buried at 12-inch depth in the center of each bed to irrigate the rows in the plot. The flow rate for the drip tape was 0.34 gal/min/100 ft at 8 psi with emitters spaced 16 inches apart, resulting in a water application rate of 0.066 inch/hour.

On October 30, 2012, seed of 11 species (Table 1) was planted in either 15-inch or 30-inch rows using a custom-made small-plot grain drill with disc openers. All seed was planted on the soil surface at 20-30 seeds/ft of row. After planting, sawdust was applied in a narrow band over the seed row at 0.26 oz/ft of row (558 lb/acre). Following planting and sawdust application, the beds were covered with row cover (N-sulate, DeWitt Co., Inc., Sikeston, MO), which covered four rows (two beds) and was applied with a mechanical plastic mulch layer. *Cymopterus bipinnatus* was planted on November 25, 2009, and *Crepis intermedia* was planted on November 28, 2011 as previously described using similar methods.

Weeds were controlled by hand-weeding as necessary.

Starting in March following fall planting, the row cover was removed. Immediately following the removal of the row cover, bird netting was placed over the seedlings on no. 9 galvanized wire hoops to prevent bird feeding on young seedlings and new shoots. During seedling emergence, wild bird seed was placed several hundred feet from the trial to attract quail away from the trials. Bird netting was removed in early May. Bird netting was applied and removed each spring.

On April 13, 2012, 50 lb nitrogen (N)/acre, 10 lb phosphorus (P)/acre, and 0.3 lb iron (Fe)/acre was applied to all plots of *Cymopterus bipinnatus* and *C. intermedia* as liquid fertilizer injected through the drip tape.

### Cultural practices in 2013

On July 26, all plots of *Machaeranthera canescens* were sprayed with Capture® at 19 oz/acre (0.3 lb ai/acre) for aphid control. On October 31, seed of *Phacelia linearis* was planted as previously described.

Due to poor stand, seed of *Chaenactis douglasii* was replanted on November 1, as previously described. Stand of *Nicotiana attenuata* was extremely poor and seed was unavailable for replanting.

### Cultural practices in 2014

Stand of *Chaenactis douglasii*, which was replanted in the fall of 2013, was poor and did not allow evaluation of irrigation responses.

On November 11, *Phacelia linearis*, *Nicotiana attenuata*, and *Thelypodium milleflorum* were replanted as previously described. Lengths of row with missing stand in plots of *Chaenactis douglasii* were replanted by hand and row cover was not applied to the replanting.

### Cultural practices in 2015

On November 2, *Nicotiana attenuata* and *Enceliopsis nudicaulis* were replanted as previously described. Before planting, the ground was not tilled, only cultipacked. On November 5,

*Phacelia linearis*, *Chaenactis douglasii*, *Achillea millefolium*, and *Ipomopsis aggregata* were replanted as previously described.

### **Cultural practices in 2016**

On November 22, *Nicotiana attenuata*, *Phacelia linearis*, and *Thelypodium milleflorum* were replanted as previously described.

### **Cultural practices in 2017**

On October 19, Prowl<sup>®</sup> H<sub>2</sub>O at 2 pt/acre was broadcast on all plots of *Enceliopsis nudicaulis*, *Crepis intermedia*, and *Thelypodium milleflorum* for weed control. On November 8, *Ipomopsis aggregata* was replanted. On November 14, *Nicotiana attenuata* was replanted.

### **Cultural practices in 2018**

Liquid fertilizer containing 0.3 lb Fe/acre was injected using a brief pulse of water through the drip irrigation system to all plots of *Crepis intermedia*, *Thelypodium milleflorum*, and *Cymopterus bipinnatus* on May 3.

### **Irrigation for seed production**

In March 2010 for *Cymopterus bipinnatus*, and March 2013 for the other species, the planted strip of each wildflower species was divided into 12 30-ft-long plots. Each plot contained four rows of each species. The experimental design for each species was a randomized complete block with four replicates. The three treatments were a non-irrigated check, 1 inch of water per irrigation, and 2 inches of water per irrigation. Each treatment received four irrigations that were applied approximately every 2 weeks starting at bud formation and flowering. The amount of water applied to each treatment was calculated by the length of time necessary to deliver 1 or 2 inches through the drip system. Irrigations were regulated with a controller and solenoid valves.

The drip-irrigation system was designed to allow separate irrigation of each species due to different timings of flowering and seed formation. All species were irrigated separately except the two *Phacelia* spp. and the two *Ligusticum* spp. Flowering, irrigation, and harvest dates were recorded (Table 2) except for *Nicotiana attenuata*, which did not germinate in 2014 and the *Ligusticum* spp., which did not flower.

### **Harvest**

All species were harvested manually in 2013. Due to a long flowering duration, seed of *Enceliopsis nudicaulis*, *Chaenactis douglasii*, and *Crepis intermedia* required multiple harvests. Seed of *E. nudicaulis* was harvested manually once a week. Seed of *Chaenactis douglasii* and *Crepis intermedia* was harvested weekly with a leaf blower in vacuum mode. In 2016, the duration of flowering for *C. intermedia* was much shorter and uniform in timing between irrigation treatments. In 2016-2018, seed of *C. intermedia* was harvested by mowing and bagging just prior to the seed heads opening. In 2016 a seed sample from each plot of *C. intermedia* was cleaned manually to determine the proportion of pure seed. A sample of light yellow (immature) seed and dark brown (mature) seed of *C. intermedia* was analyzed for viability (tetrazolium). In 2016, seed of *Chaenactis douglasii* was harvested manually once a week.

*Machaeranthera canescens* seed was harvested by cutting and windrowing the plants. After drying for 2 days the *M. canescens* plants were beaten on plastic tubs to separate the seed heads from the stalks. *Phacelia hastata* was harvested with a small-plot combine in 2014 and 2015. In

2016 and 2017, *P. hastata* was harvested manually due to the low stature of the plants. *Heliomeris multiflora* was harvested with a small plot combine in 2015 and 2016. The duration of flowering for *H. multiflora* tends to increase with increasing irrigation. In 2013 and 2014, the duration of flowering in the wetter plots of *H. multiflora* was much longer than in the drier plots, making a single mechanical harvest unfeasible. In 2015, the duration of flowering in the wetter plots of *H. multiflora* was shorter, enabling mechanical harvest. In 2016, plots of the driest treatment were harvested manually before the other plots, which were harvested mechanically on July 8. All plots of *H. multiflora* were harvested with a small plot combine in 2017.

Seed of all species was cleaned manually.

## Statistical analysis

Seed yield means were compared by analysis of variance and by linear and quadratic regression. Seed yield ( $y$ ) in response to irrigation or irrigation plus precipitation ( $x$ , inches/season) was estimated by the equation  $y = a + b \cdot x + c \cdot x^2$ . For the quadratic equations, the amount of irrigation ( $x'$ ) that resulted in maximum yield ( $y'$ ) was calculated using the formula  $x' = -b/2c$ , where  $a$  is the intercept,  $b$  is the linear parameter, and  $c$  is the quadratic parameter. For the linear regressions, the seed yield responses to irrigation were based on the actual greatest amount of water applied plus precipitation and the measured average seed yield.

## Results and Discussion

Precipitation in the winter and spring in 2013 was lower and in 2017 was higher than the 5-year average (Table 3). In 2018, precipitation in the fall, winter, and spring was lower than average. Precipitation in the other years was close to the average. The accumulation of growing degree-days (50-86°F) was higher than average in 2013-2016 and in 2018 (Table 3). In 2017, the accumulation of growing degree-days was close to the average.

***Achillea millefolium***. Flowering and seed production in 2016, the first year after fall planting, was minimal. Seed yields of *A. millefolium* showed a quadratic response to irrigation in 2017 and 2018 with maximum seed yields of 235 lb/acre and 57 lb/acre at 6.2 and 7.9 inches of water applied in 2017 and 2018, respectively (Tables 4 and 5).

***Chaenactis douglasii***. Stands of *C. douglasii* were poor in 2013 and 2014 and did not permit evaluation of irrigation responses. Replanting in the fall of 2013-2015 was necessary to establish an adequate stand of *C. douglasii*, allowing evaluations of irrigation responses in 2015-2018. *Chaenactis douglasii* seed yields did not respond to irrigation in 2015-2017. In 2018, seed yield showed a quadratic response to irrigation with maximum seed yields of 85 lb/acre at 6.3 inches of water applied. Highest seed yields averaged 225 lb/acre over the 4 years.

***Crepis intermedia***. *Crepis intermedia* flowered and produced seed for the first time in 2015, the third year after fall planting in 2011. The uniform and short flowering of *C. intermedia* in 2016 allowed the seed from all plots to be harvested once. A single mechanical harvest is more efficient, but some of the seed could be immature because harvest needed to occur just before seed heads opened. In 2016, 77% of the seed harvested was mature and had a viability of 57%. The other 23% of the harvested seed was immature and had a viability of 5%. This suggests that a single harvest as conducted in this trial resulted in adequate seed quality. *Crepis intermedia* seed yields increased with increasing irrigation rate up to the highest rate of 8 inches in 2015. In 2016 and 2017, seed yields of *C. intermedia* did not respond to irrigation. In 2018, seed yield

showed a quadratic response to irrigation with maximum seed yields of 151 lb/acre at 4 inches of water applied. Seed yields increased each year from 2015 to 2017 with highest seed yields of 302 lb/acre in 2017. Seed yields were lower in 2018.

***Cymopterus bipinnatus***. *Cymopterus bipinnatus* did not flower in either 2010 or 2011 and flowered very little in 2012. *Cymopterus bipinnatus* seed yields did not respond to irrigation in 2013 and 2016. In 2014, seed yields increased with increasing irrigation rate up to the highest rate of 8 inches. In 2015 and 2018, seed yields showed a quadratic response to irrigation with maximum seed yields at 4.2 and 4.8 inches of water applied in 2015 and 2018, respectively. In 2017, seed yields were highest with no irrigation. Averaged over the 6 years, seed yields were estimated to be highest with 5.2 inches of total applied irrigation water yielding 975 lb/acre of seed.

***Enceliopsis nudicaulis***. *Enceliopsis nudicaulis* seed yield was very low and did not respond to irrigation in 2013. In 2014, seed yield showed a quadratic response to irrigation with a maximum seed yield at 5.4 inches of water applied. Extensive die-off of *E. nudicaulis* occurred over the winter of 2014-2015 and was more severe in the plots receiving the highest amount of irrigation. Seed yields of *E. nudicaulis* were substantially reduced in 2015 and were highest without irrigation. In 2016, seed yield showed a quadratic response to irrigation with a maximum seed yield at 5.8 inches of water applied. In 2017, seed yields were highest without irrigation. Seed yields did not respond to irrigation in 2018. The replanting done in the fall of 2015 was successful, but stands continue to decline, especially in the irrigated plots. Highest seed yields averaged 26 lb/acre over the 4 years.

***Heliomeris multiflora***. *Heliomeris multiflora* seed yield increased with increasing irrigation rate up to the highest rate of 8 inches in 2013-2015; *H. multiflora* seed yield did not respond to irrigation in 2016 and 2017. In 2018, seed yields showed a quadratic response to irrigation with a maximum seed yield at 3.7 inches of water applied. Highest seed yields averaged 130 lb/acre over the 6 years.

***Ipomopsis aggregata***. *Ipomopsis aggregata* flowered very little in 2013, then flowered and set seed in 2014. The stand of *I. aggregata* died over the winter of 2014-2015, which indicated a biennial growth habit. *Ipomopsis aggregata* seed yields were highest with 4 inches of water applied in 2014 and 2017. Highest seed yields averaged 262 lb/acre over the 2 years.

***Machaeranthera canescens***. *Machaeranthera canescens* seed yields showed a quadratic response to irrigation with a maximum seed yield at 2.4 inches of water applied in 2013. In 2014, 2015, 2018, and averaged over the 3 years, seed yields of *M. canescens* did not respond to irrigation. Highest seed yields averaged 700 lb/acre over the 4 years. Partial die-off of *M. canescens* over the winter of 2015-2016 resulted in stand too uneven for an irrigation trial in 2016 and 2017. Natural reseeding occurred over the winter of 2016-2017, but the young plants did not flower in 2017.

***Nicotiana attenuata***. Stand establishment has been difficult with only one fall planting (2015) out of six resulting in adequate stand for the irrigation trial. Seed yields of *N. attenuata* showed a quadratic response to irrigation in 2016 with a maximum seed yield of 151 lb/acre at 4.6 inches of water applied

***Phacelia hastata***. Irrigation responses for *P. hastata* were evaluated for two sets of plots: the 6-year-old stand planted in 2012 and a new stand originating in 2015 from volunteer seed. *Phacelia hastata* (planted in the fall of 2012) seed yields showed a quadratic response to

irrigation with a maximum seed yield at 5.3, 7.6, and 5.2 inches of water applied in 2013, 2014, and 2018, respectively. In 2015, seed yield of *P. hastata* did not respond to irrigation, possibly due to loss of stand in this weak perennial. The original stand of *P. hastata*, planted in the fall of 2012, was extremely poor in 2016 and seed was not harvested. Stand regenerated from natural reseeding in 2017, but seed was not produced.

Seed yields of *P. hastata* (started in the fall of 2014) increased with increasing irrigation rate up to the highest rate of 8 inches in 2015. In 2016 and 2018, seed yields of *P. hastata* showed a quadratic response to irrigation with a maximum seed yield at 4 and 6.8 inches of water applied in 2016 and 2018, respectively. In 2017, seed yields of *P. hastata* did not respond to irrigation. Averaged over the 4 years, seed yields of *P. hastata* showed a quadratic response to irrigation with a maximum seed yield of 162 lb/acre and 83 lb/acre at 6.2 and 5.5 inches of water applied for the 2012 and 2014 stands, respectively. The two stands of *P. hastata* showed a pattern of increased seed yields in the second year, a decline in the third year, and an increase in the fourth year.

***Phacelia linearis*.** Seed yields of *P. linearis* showed a quadratic response to irrigation in 2013 with a maximum seed yield at 6.2 inches of water applied. In 2014, seed yields of *P. linearis* did not respond to irrigation. Highest seed yields averaged 240 lb/acre over the 2 years.

The replanting of *P. linearis* in the fall of 2014 and 2015 did not result in adequate stands. *Phacelia linearis* was replanted in the fall of 2016 in a different location in the field but stand in the spring of 2017 was extremely poor.

***Thelypodium milleflorum*.** Seed yield of *T. milleflorum* did not respond to irrigation in any of the 3 years of seed production (Tables 4 and 5). Highest seed yields averaged 152 lb/acre over the 3 years.

Stands of *Ligusticum porteri* and *L. canbyi* were poor and uneven and did not permit evaluation of irrigation responses.

## Acknowledgements

This project was funded by the U.S. Forest Service Great Basin Native Plant Project, U.S. Bureau of Land Management, Oregon State University, Malheur County Education Service District, and supported by Formula Grant nos. 2018-31100-06041 and 2018-31200-06041 from the USDA National Institute of Food and Agriculture.

Table 2. Native wildflower flowering, irrigation, and seed harvest dates by species. Malheur Experiment Station, Oregon State University, Ontario, OR, 2013-2018. Table 2 is continued on the next page.

Year	Flowering dates			Irrigation dates		Harvest
	Start	Peak	End	Start	End	
<b><i>Achillea millefolium</i>, common yarrow</b>						
2017	26-Apr	7-Jun	12-Jul	2-May	20-Jun	26-Jul
2018	30-Apr	13-Jun	30-Jun	16-May	27-Jun	3-Aug
<b><i>Chaenactis douglasii</i>, Douglas' dustymaiden</b>						
2013	23-May	30-Jun	15-Jul	22-May	3-Jul	2-Jul, 22-Jul
2014	20-May		15-Jul	13-May	24-Jun	poor stand
2015	5-May		10-Jul	5-May	17-Jun	weekly, 8-Jun to 15-Jul
2016	23-May		22-Jul	23-May	8-Jul	weekly, 17-Jun to 7-Jul
2017	25-May	7-Jun	19-Jul	9-May	20-Jun	weekly, 16-Jun to 6-Jul
2018	10-May	13-Jun	10-Jul	16-May	27-Jun	weekly, 13-Jun to 15-Jul
<b><i>Machaeranthera canescens</i>, hoary tansyaster</b>						
2013	13-Aug		1-Oct	17-Jul	28-Aug	2-Oct
2014	20-Aug	17-Sep	5-Oct	22-Jul	2-Sep	6-Oct
2015	10-Aug	17-Sep	1-Oct	11-Aug	22-Sep	6-Oct, 15-Oct
2016	17-Aug	20-Sep	10-Oct			partial winter die-off
2017	29-Aug		20-Oct			
2018	20-Aug		22-Oct	23-Aug	20-Sep	22-Oct
<b><i>Phacelia hastata</i>, silverleaf phacelia</b>						
2013	17-May		30-Jul	22-May	3-Jul	30-Jul (0 in), 7-Aug, 19-Aug (8 in)
2014	5-May		10-Jul	29-Apr	10-Jun	14-Jul
2015 (1st year)	28-Apr	26-May	7-Aug	20-May	30-Jun	6-Aug
2015 (3rd year)	28-Apr	26-May	7-Aug	29-Apr	10-Jun	7-Jul (0 in), 21-Jul (4, 8 in)
2016	28-Apr		17-Jun	27-Apr	7-Jun	23-Jun
2017	8-May	7-Jun		2-May	20-Jun	25-Jul
2018	6-May		20-Jun	16-May	27-Jun	27-Jun
<b><i>Phacelia linearis</i>, threadleaf phacelia</b>						
2013	3-May	16-May	15-Jun	2-May	12-Jun	2-Jul
2014	5-May	4-Jun	1-Jul	1-May	10-Jun	7-Jul
2015			winter die-off			
<b><i>Enceliopsis nudicaulis</i>, nakedstem sunray</b>						
2013	30-Jun		15-Sep	3-Jul	14-Aug	weekly, 8-Aug to 30-Aug
2014	5-May	1-Jul	30-Jul	6-May	17-Jun	weekly, 14-Jul to 30-Aug
2015	28-Apr	13-May	5-Aug	29-Apr	10-Jun	weekly, 2-Jun to 15-Aug
2016	20-Apr		30-Jul	3-May	14-Jun	weekly, 27-Apr to 29-Jul
2017	11-May	7-Jun	20-Aug	23-May	6-Jul	weekly, 4-Jun to 15-Aug
2018	30-Apr	26-Jun	30-Jul	16-May	27-Jun	weekly, 27-Apr to 27-Jul

Table 2. (Continued.) Native wildflower flowering, irrigation, and seed harvest dates by species. Malheur Experiment Station, Oregon State University, Ontario, OR, 2013-2018.

Year	Flowering dates			Irrigation dates		Harvest
	Start	Peak	End	Start	End	
<b><i>Heliomeris multiflora</i>, showy goldeneye</b>						
2013	15-Jul		30-Aug	5-Jun	17-Jun	8-Aug, 15-Aug, 28-Aug
2014	20-May	20-Jun	30-Aug	13-May	24-Jun	weekly, 15-Jul to 15-Aug
2015	5-May	26-May	10-Jul	5-May	17-Jun	13-Jul
2016	5-May	15-Jun	30-Sep	9-May	22-Jun	8-Jul
2017	12-May	7-Jun	30-Jul	9-May	20-Jun	17-Jul
2018	12-May	13-Jun	20-Jul	16-May	27-Jun	1-Aug
<b><i>Cymopterus bipinnatus</i>, Hayden's cymopterus</b>						
2013	5-Apr		15-May	12-Apr	22-May	10-Jun
2014	7-Apr		29-Apr	7-Apr	20-May	16-Jun
2015	25-Mar		24-Apr	1-Apr	13-May	8-Jun
2016	15-Mar		25-Apr	31-Mar	9-May	7-Jun
2017	27-Mar		1-May	19-Apr	6-Jun	16-Jun
2018	15-Mar		3-May	18-Apr	30-May	5-Jun
<b><i>Ipomopsis aggregata</i>, scarlet gilia</b>						
2013	31-Jul	very little flowering		31-Jul	11-Sep	
2014	22-Apr	13-May	30-Jul	23-Apr	3-Jun	20-Jun
2015		winter die-off				
2016		No flowering		7-Jun	22-Jul	
2017	1-May	15-May	27-Jun	2-May	20-Jun	23-Jun
2018		No flowering		16-May	27-Jun	
<b><i>Thelypodium milleflorum</i>, manyflower thelypody</b>						
2013		No flowering				
2014	22-Apr	5-May	10-Jun	23-Apr	3-Jun	2-Jul
2015		No flowering				
2016	11-Apr	6-May	8-Jun	11-Apr	23-May	21-Jun
2017		No flowering				
2018	27-Apr	10-May	10-Jun	3-May	13-Jun	18-Jun
<b><i>Crepis intermedia</i>, limestone hawkbeard</b>						
2015	28-Apr	5-May	1-Jun	21-Apr	3-Jun	weekly, 1-Jun to 2-Jul
2016	29-Apr		25-May	27-Apr	7-Jun	26-May
2017	15-May		7-Jun	9-May	20-Jun	8-Jun
2018	3-May		25-May	3-May	13-Jun	31-May
<b><i>Nicotiana attenuata</i>, coyote tobacco</b>						
2016	16-May		31-Jul	16-May	22-Jun	weekly, 21-Jun to 29-Jul
2017	1-May		15-Aug			



Table 3. Precipitation and growing degree-days at the Malheur Experiment Station, Ontario, OR, 2013-2018.

Year	Precipitation (inch)			Growing degree-days (50-86°F)
	Spring	Winter + spring	Fall + winter + spring	Jan–Jun
2013	0.9	2.4	5.3	1319
2014	1.7	5.1	8.1	1333
2015	3.2	5.9	10.4	1610
2016	2.2	5.0	10.1	1458
2017	4.0	9.7	12.7	1196
2018	1.9	4.9	5.8	1342
5-year average:	2.4	5.6	9.3	25-year average: 1207

Table 4. Native wildflower seed yield (lb/acre) in response to season-long irrigation rate (inches). Malheur Experiment Station, Oregon State University, Ontario, OR, 2013-2018. Table 4 is continued on the next page.

Species	Year	Irrigation rate			LSD (0.05)
		0 inches	4 inches	8 inches	
		----- lb/acre -----			
<i>Achillea millefolium</i>	2017	59.2	213.3	220.4	99.8
	2018	7.3	45.1	57.1	NS
	Average	23.2	93.2	94.0	46.0
<i>Chaenactis douglasii</i>	2015	132.1	137.6	183.3	NS <sup>a</sup>
	2016	29.1	16.0	27.2	NS
	2017	707.1	711.1	627.3	NS
	2018	7.9	74.7	79.6	12.5
	Average	208.3	213.7	225.4	NS
<i>Crepis intermedia</i>	2015	68.6	55.5	166.3	63.2
	2016	83.6	87.0	77.8	NS
	2017	301.5	268.1	287.1	NS
	2018	98.3	151.0	100.2	16.1
	Average	138.0	140.4	155.5	NS
<i>Cymopterus bipinnatus</i>	2013	194.2	274.5	350.6	NS
	2014	1236.2	1934.0	2768.5	844.7
	2015	312.3	749.0	374.9	240.7
	2016	1501.4	2120.6	1799.0	546.6 <sup>b</sup>
	2017	245.4	178.6	95.8	NS
	2018	87.0	149.5	122.5	15.8
	Average	618.8	956.4	868.2	153.2
<i>Enceliopsis nudicaulis</i>	2013	2.3	6.8	5.9	NS
	2014	1.5	34.6	29.1	20.7
	2015	15.7	3.2	4.4	7.3
	2016	10.5	47.6	45.9	34.9
	2017	105.0	43.2	25.0	59.6
	2018	20.2	20.5	20.1	NS
	Average	25.9	26.2	21.7	NS

Table 4. (Continued.) Native wildflower seed yield (lb/acre) in response to season-long irrigation rate (inches). Malheur Experiment Station, Oregon State University, Ontario, OR, 2013-2018.

Species	Year	Irrigation rate			LSD (0.05)
		0 inches	4 inches	8 inches	
		----- lb/acre -----			
<i>Helioomeris multiflora</i>	2013	28.7	57.6	96.9	NS
	2014	154.6	200.9	271.7	107.3 <sup>b</sup>
	2015	81.7	115.6	188.2	58.2
	2016	92.3	89.2	98.0	NS
	2017	87.8	75.9	89.9	NS
	2018	44.5	73.9	34.3	23.4
	Average	84.0	101.4	129.8	24.7
<i>Ipomopsis aggregata</i>	2014	47.1	60.9	63.6	9.0
	2017	241.0	315.8	188.8	74.5
	Average	180.3	261.7	145.1	97.2
<i>Machaeranthera canescens</i>	2013	206.1	215	124.3	73.6
	2014	946.1	1210.2	1026.3	NS
	2015	304.1	402.6	459.1	NS
	2018	330.3	426.3	380.6	NS
	Average	586.1	701.6	634.3	NS
<i>Nicotiana attenuata</i>	2016	49.4	151.0	95.8	81.4
<i>Phacelia hastata</i> (planted fall 2012)	2013	35.3	102.7	91.2	35.7
	2014	87.7	305.7	366.4	130.3
	2015	78.8	79.3	65.0	NS
	2018	32.8	108.6	89.6	59.4
	Average	58.6	149.1	153.0	37.0
<i>Phacelia hastata</i> (planted fall 2014)	2015	0.0	21.4	50.4	13.7
	2016	82.5	125.2	83.1	26.8
	2017	20.3	23.2	23.2	NS
	2018	57.1	128.5	140.2	68.3
	Average	40.0	79.6	74.2	22.0
<i>Phacelia linearis</i>	2013	121.4	306.2	314.2	96
	2014	131.9	172.9	127.2	NS
	Average	126.7	239.5	220.7	87.2
<i>Thelypodium milleflorum</i>	2014	200.5	246.2	205.6	NS
	2016	121.9	110.0	63.3	NS
	2018	61.4	61.4	64.1	NS
	Average	131.5	151.7	117.0	NS

<sup>a</sup>Not significant. <sup>b</sup>LSD (0.10).

Table 5. Regression analysis for native wildflower seed yield (y) in response to irrigation (x) (inches/season) using the equation  $y = a + b \cdot x + c \cdot x^2$ . For the quadratic equations, the amount of irrigation that resulted in maximum yield was calculated using the formula:  $-b/2c$ , where b is the linear parameter and c is the quadratic parameter. Malheur Experiment Station, Oregon State University, Ontario, OR, 2013-2018. Table 5 is continued on the next page.

Species	Year	intercept	linear	quadratic	$R^2$	$P$	Maximum yield lb/acre	Water applied for maximum yield inches/season
<b><i>Achillea millefolium</i></b>	2017	59.2	56.9	-4.6	0.75	0.01	235.4	6.2
	2018	7.3	12.7	-0.8	0.49	0.1	57.1	7.9
	Average	23.2	26.2	-2.2	0.72	0.01	102.3	6.0
	<hr/>							
<b><i>Chaenactis douglasii</i></b>	2015	125.4	6.4		0.08	NS <sup>a</sup>		
	2016	25.1	-0.2		0.01	NS		
	2017	707.1	12.0	-2.7	0.09	NS		
	2018	7.9	24.4	-1.9	0.99	0.001	85.1	6.3
	Average	207.2	2.1		0.04	NS		
<hr/>								
<b><i>Crepis intermedia</i></b>	2015	49.0	11.4		0.31	0.10	176.6	8.0
	2016	83.6	2.4	-0.4	0.07	NS		
	2017	292.8	-1.8		0.01	NS		
	2018	98.3	26.1	-3.2	0.41	0.10	151.0	4.0
	Average	135.5	2.4		0.07	NS		
<hr/>								
<b><i>Cymopterus bipinnatus</i></b>	2013	194.9	19.6		0.07	NS		
	2014	1214.6	190.6		0.41	0.05	2739.4	8.0
	2015	312.3	210.5	-25.3	0.46	0.10	749.6	4.2
	2016	1501.4	272.4	-29.4	0.34	NS		
	2017	308.1	-24.4		0.38	0.10	308.1	0.0
	2018	87.0	26.8	-2.8	0.60	0.05	151.3	4.8
	Average	618.8	137.6	-13.3	0.52	0.05	974.6	5.2
	<hr/>							
<b><i>Enceliopsis nudicaulis</i></b>	2013	3.1	0.4		0.16	NS		
	2014	1.5	13.1	-1.2	0.6	0.05	37.3	5.5
	2015	13.4	-1.4		0.29	0.10	13.4	0.0
	2016	10.5	14.1	-1.2	0.57	0.05	51.6	5.8
	2017	99.1	-10.0		0.44	0.05	99.1	0.0
	2018	20.3	0.0		0.01	NS		
	Average	25.9	0.7	-0.1	0.04	NS		

<sup>a</sup>Not significant. There was no statistically significant trend in seed yield in response to amount of irrigation.

Table 5. (Continued.) Regression analysis for native wildflower seed yield (y) in response to irrigation (x) (inches/season) using the equation  $y = a + bx + cx^2$ . For the quadratic equations, the amount of irrigation that resulted in maximum yield was calculated using the formula:  $-b/2c$ , where b is the linear parameter and c is the quadratic parameter. Malheur Experiment Station, Oregon State University, Ontario, OR, 2013-2018.

Species	Year	intercept		quadratic	$R^2$	P	Maximum	Water applied
		linear	linear				yield	for maximum
							lb/acre	inches/season
<b><i>Helioomeris multiflora</i></b>	2013	27	8.5		0.38	0.05	95.0	8.0
	2014	150.5	14.6		0.27	0.10	267.3	8.0
	2015	75.2	13.3		0.48	0.05	181.8	8.0
	2016	90.7	0.7		0.01	NS		
	2017	83.5	0.3		0.01	NS		
	2018	44.5	16.0	-2.2	0.72	0.01	74.1	3.7
	Average	82.1	5.7		0.44	0.05	128.0	8.0
<b><i>Ipomopsis aggregata</i></b>	2014	48.5	2.1		0.23	NS		
	2017	241.0	43.9	-6.3	0.52	0.05	317.5	3.5
	Average	180.3	45.1	-6.2	0.24	NS		
<b><i>Machaeranthera canescens</i></b>	2013	206.1	14.7	-3.1	0.54	0.05	223.5	2.4
	2014	946.1	122	-14	0.13	NS		
	2015	311.1	19.4		0.02	NS		
	2018	330.3	41.7	-4.4	0.03	NS		
	Average	586.1	51.7	-5.7	0.09	NS		
<b><i>Nicotiana attenuata</i></b>	2016	49.4	45.0	-4.9	0.50	0.05	152.7	4.6
<b><i>Phacelia hastata</i></b> (planted fall 2012)	2013	35.3	26.7	-2.5	0.66	0.01	106.6	5.3
	2014	87.7	74.2	-4.9	0.76	0.01	368.6	7.6
	2015	78.8	2.0	-0.5	0.04	NS		
	2018	32.8	30.8	-3.0	0.49	0.05	112.9	5.2
	Average	58.6	33.4	-2.7	0.84	0.001	162.0	6.2
<b><i>Phacelia hastata</i></b> (planted fall 2014)	2015	-1.3	6.3		0.88	0.001	49.2	8.0
	2016	82.5	21.3	-2.6	0.72	0.01	125.2	4.0
	2017	20.3	1.1	-0.1	0.04	NS		
	2018	57.1	25.3	-1.9	0.57	0.05	143.0	6.8
	Average	40.0	15.5	-1.4	0.69	0.01	82.8	5.5
<b><i>Phacelia linearis</i></b>	2013	121.4	68.3	-5.5	0.69	0.01	333.4	6.2
	2014	131.9	21.1	-2.7	0.11	NS		
	Average	126.7	44.7	-4.1	0.48	0.1	248.5	5.5
<b><i>Thelypodium milleflorum</i></b>	2014	200.5	22.2	-2.7	0.12	NS		
	2016	121.9	1.4	-1.1	0.35	NS		
	2018	61.0	0.3		0.01	NS		
	Average	131.5	11.9	-1.7	0.16	NS		

<sup>a</sup>Not significant. There was no statistically significant trend in seed yield in response to amount of irrigation.